arXiv:1105.1781v2 [hep-ph] 9 Jun 2011

Dark Matter from Minimal Flavor Violation

Brian Batell,[1, *] Josef Pradler,[1, †] and Michael Spannowsky[2, ‡]
[1] *Perimeter Institute for Theoretical Physics, Waterloo, ON, N2L 2Y5, Canada*
[2] *Institute of Theoretical Science, University of Oregon, Eugene, OR 97403-5203, USA*

We consider theories of flavored dark matter, in which the dark matter particle is part of a multiplet transforming nontrivially under the flavor group of the Standard Model in a manner consistent with the principle of Minimal Flavor Violation (MFV). MFV automatically leads to the stability of the lightest state for a large number of flavor multiplets. If neutral, this particle is an excellent dark matter candidate. Furthermore, MFV implies specific patterns of mass splittings among the flavors of dark matter and governs the structure of the couplings between dark matter and ordinary particles, leading to a rich and predictive cosmology and phenomenology. We present an illustrative phenomenological study of an effective theory of a flavor $SU(3)_Q$ triplet, gauge singlet scalar.

The observational evidence for dark matter (DM) provides one of the few solid empirical motivations for new particle physics [1]. Indeed, the simplest explanation of DM is a new massive particle that is stable on cosmological time scales and a thermal relic of the Big Bang. Many scenarios for physics beyond the Standard Model (SM) include candidates for the DM particle, with a popular example being the neutralino in theories with weak scale supersymmetry (SUSY).

Typically, in such theories the stability of the DM particle is a consequence of a discrete Z_2 parity symmetry motivated by the need to eliminate dangerous operators allowed by SM gauge invariance. For example, in the Minimal Supersymmetric Standard Model (MSSM) renormalizable terms in the superpotential can violate baryon and lepton number and thus lead to rapid proton decay. R-parity conservation is often invoked to eliminate such operators, with the welcome side effect of rendering the lightest supersymmetric particle stable and therefore a DM candidate.

It is remarkable that certain phenomenological constraints such as proton decay may be overcome by invoking discrete symmetries, which in turn lead to stable DM candidates. However, our detailed knowledge of the flavor sector of the SM typically requires a highly non-trivial structure in the couplings of new particles to the quarks and leptons—one that is not easily achieved by imposing discrete symmetries. For example, R-parity does not prevent large flavor changing neutral currents (FCNCs) in the MSSM, and additional assumptions about the structure of SUSY breaking must be made. In this regard the principle of Minimal Flavor Violation (MFV) is well motivated [2, 3]. One postulates that the global flavor symmetry of the SM is broken only by the Yukawa matrices in the presence of new particles. This provides a built-in protection mechanism against large flavor violating processes. We shall demonstrate that, in analogy with the discrete symmetries discussed above, the MFV principle can lead to stable neutral particles and excellent DM candidates.

In this paper we propose new theories of flavored DM. We add to the SM new matter multiplets with nontrivial transformation properties under the flavor group. We show that the MFV hypothesis implies that certain flavor representations contain one or more stable components, and we identify the general conditions for stability in these theories. Many variants of flavored DM models can then be constructed, each with a rich and predictive phenomenology.

As an example, we study a simple effective theory of DM with a flavor $SU(3)_Q$ triplet, gauge singlet scalar field, which provides a representative survey of various aspects of the cosmology and phenomenology in the flavored DM scenario. MFV predicts patterns in the mass splittings of the DM flavors governed by the SM Yukawa couplings. Flavor is intertwined with the physics of DM, as the Yukawa couplings and CKM matrix play a role in the thermal freezeout, direct and indirect signatures, and collider signals. In the scenario studied here, the heavy flavors in the DM multiplet decay to lighter flavors along with SM particles, leading to striking signatures at collider experiments. In other models, it is possible that the heavy flavors are stable on cosmological time scales, in which case several species comprise the cosmic DM.

We note that several related studies exist in the literature. As an alternative to R-parity, Ref. [4] considers MFV as a protection mechanism in the MSSM, although in this scenario the LSP is not stable on cosmological times scales and is therefore not a DM candidate. Supersymmetric DM tied to the neutrino flavor structure has been considered in [5] while Ref. [6] investigated theories of flavored DM, though both of these proposals are outside the MFV framework. Lastly, and in a more general context, models of flavor employing *e.g.* discrete symmetries [7, 8] or horizontal gauge symmetries [9] have also found their application in the DM problem.

Flavor symmetries and dark matter. We begin by reviewing the MFV hypothesis, which provides a mechanism based on flavor symmetries by which new light particles may be consistent with precision flavor observables. The quark sector of SM exhibits a large global flavor symmetry $G_q = SU(3)_Q \times SU(3)_{u_R} \times SU(3)_{d_R}$, which is broken only by Yukawa interactions. When considering theories of physics beyond the SM, the MFV principle dictates that new particles also respect G_q and the only sources of flavor breaking arise from insertions of the SM Yukawa matrices Y_u and Y_d [2, 3]. Formally, this is achieved by promoting the Yukawa matrices to spurion fields which transform nontrivially under G_q. The SM Yukawa interactions are written as

$$-\mathcal{L}_Y \supset \bar{Q} Y_d d_R H + \bar{Q} Y_u u_R H^\dagger + \text{h.c.}, \tag{1}$$

so that the Yukawa spurions transform as $Y_u \sim (\mathbf{3}, \bar{\mathbf{3}}, \mathbf{1})$, $Y_d \sim (\mathbf{3}, \mathbf{1}, \bar{\mathbf{3}})$ under G_q.

A number of studies considering new MFV matter content exist in the literature [10–15]. In all of these works, the G_q representations of the new particles have been chosen such that renormalizable tree-level couplings to the SM fermions are allowed. An immediate consequence of this is that such particles are unstable and decay to SM quarks or leptons. We now investigate the generality of this conclusion, and show in particular that MFV implies stability for particles in certain representations of G_q. With further assumptions about their electroweak quantum numbers, these stable particles become perfectly viable DM candidates.

Let χ be a matter multiplet that transforms nontrivially under flavor but is color-neutral, *i.e.* a singlet under $SU(3)_c$. The most general operator that induces the decay of χ then reads

$$\mathcal{O}_{\rm decay} = \chi\, \underbrace{Q\ldots}_{A}\underbrace{\bar{Q}\ldots}_{B}\underbrace{u_R\ldots}_{C}\underbrace{\bar{u}_R\ldots}_{D}\underbrace{d_R\ldots}_{E}\underbrace{\bar{d}_R\ldots}_{F} \times \underbrace{Y_u\ldots}_{G}\underbrace{Y_u^\dagger\ldots}_{H}\underbrace{Y_d\ldots}_{I}\underbrace{Y_d^\dagger\ldots}_{J}\,\mathcal{O}_{\rm weak}, \tag{2}$$

where A is the number of Q fields, B is the number of $\bar{Q}$ fields, etc. $\mathcal{O}_{\rm weak}$ is a potential electroweak operator so that (2) is rendered invariant under $SU(2)_L \times U(1)_Y$. The flavor and color indices of the remaining fields are to be contracted by use of the $SU(3)$ group invariant tensors δ and ϵ so that (2) becomes a color and flavor singlet. This, however, can only be achieved if the triality t_i of each $SU(3)_i$ tensor product $(p, q)_i$ with p factors of $\mathbf{3}_i$ and q factors of $\bar{\mathbf{3}}_i$, vanishes for the operator $\mathcal{O}_{\rm decay}$:

$$t_i \equiv (p-q)_i \bmod 3 = 0, \quad i = c,\, Q,\, u_R,\, d_R. \tag{3}$$

(n, m)	$SU(3)_Q \times SU(3)_{u_R} \times SU(3)_{d_R}$	Stable?
$(0, 0)$	$(\mathbf{1},\mathbf{1},\mathbf{1})$	
$(1, 0)$	$(\mathbf{3},\mathbf{1},\mathbf{1})$, $(\mathbf{1},\mathbf{3},\mathbf{1})$, $(\mathbf{1},\mathbf{1},\mathbf{3})$	Yes
$(0, 1)$	$(\bar{\mathbf{3}},\mathbf{1},\mathbf{1})$, $(\mathbf{1},\bar{\mathbf{3}},\mathbf{1})$, $(\mathbf{1},\mathbf{1},\bar{\mathbf{3}})$	Yes
(2,0)	$(\mathbf{6},\mathbf{1},\mathbf{1})$, $(\mathbf{1},\mathbf{6},\mathbf{1})$, $(\mathbf{1},\mathbf{1},\mathbf{6})$ $(\mathbf{3},\mathbf{3},\mathbf{1})$, $(\mathbf{3},\mathbf{1},\mathbf{3})$, $(\mathbf{1},\mathbf{3},\mathbf{3})$	Yes
(0,2)	$(\bar{\mathbf{6}},\mathbf{1},\mathbf{1})$, $(\mathbf{1},\bar{\mathbf{6}},\mathbf{1})$, $(\mathbf{1},\mathbf{1},\bar{\mathbf{6}})$ $(\bar{\mathbf{3}},\bar{\mathbf{3}},\mathbf{1})$, $(\bar{\mathbf{3}},\mathbf{1},\bar{\mathbf{3}})$, $(\mathbf{1},\bar{\mathbf{3}},\bar{\mathbf{3}})$	Yes
(1,1)	$(\mathbf{8},\mathbf{1},\mathbf{1})$, $(\mathbf{1},\mathbf{8},\mathbf{1})$, $(\mathbf{1},\mathbf{1},\mathbf{8})$ $(\mathbf{3},\bar{\mathbf{3}},\mathbf{1})$, $(\mathbf{3},\mathbf{1},\bar{\mathbf{3}})$, $(\mathbf{1},\mathbf{3},\bar{\mathbf{3}})$ $(\bar{\mathbf{3}},\mathbf{3},\mathbf{1})$, $(\bar{\mathbf{3}},\mathbf{1},\mathbf{3})$, $(\mathbf{1},\bar{\mathbf{3}},\mathbf{3})$	

TABLE I. Flavored DM candidates. Listed are the lowest-dimensional representations of G_q, organized according (n, m) where $n \equiv n_Q + n_u + n_d$, $m \equiv m_Q + m_u + m_d$. We have also indicated the representations that are stable once MFV is imposed. Depending on their electroweak quantum numbers, these multiplets may contain viable DM candidates.

On the contrary, the decay operator (2) will be forbidden if $t_i \neq 0$ for at least one i. Denoting the irreducible representation of χ under G_q as

$$\chi \sim (n_Q, m_Q)_Q \times (n_u, m_u)_{u_R} \times (n_d, m_d)_{d_R}, \tag{4}$$

where n_Q, m_Q, etc. can take values $0, 1, 2, \ldots$, the triality conditions (3) become

$$t_c = (A - B + C - D + E - F) \bmod 3 = 0, \tag{5}$$

$$t_Q = (n_Q - m_Q + A - B + G - H + I - J) \bmod 3 = 0, \tag{6}$$

$$t_{u_R} = (n_u - m_u + C - D - G + H) \bmod 3 = 0, \tag{7}$$

$$t_{d_R} = (n_d - m_d + E - F - I + J) \bmod 3 = 0. \tag{8}$$

Adding together Eqs. (6-8) and subtracting Eq. (5), we find that a necessary condition for $\mathcal{O}_{\rm decay}$ to be allowed, and thus for χ to be unstable is $(n-m) \bmod 3 = 0$, where $n \equiv n_Q + n_u + n_d$ and $m \equiv m_Q + m_u + m_d$. It follows that that $\mathcal{O}_{\rm decay}$ is forbidden and χ is stable if

$$(n - m) \bmod 3 \neq 0. \tag{9}$$

Once (9) holds, χ contains a stable component. In this regard, it is important to note that for operators with multiple fields χ, $\chi^\dagger$ which may potentially mediate a loop induced decay, the stability condition above still holds. Table I lists the lowest-dimensional representations of G_q that are stable according to the condition (9).

In order to provide a viable theory of DM, any stable state in the flavored multiplet χ must further be electrically neutral. It thus remains to specify the electroweak quantum numbers of χ. One possibility is that χ is a SM gauge singlet [16]. Alternatively, as discussed in Ref. [17], χ may be a n-plet of $SU(2)_L$ with hypercharge Y such that a component of χ is neutral, $Q = T_3 + Y = 0$, with T_3 being the diagonal $SU(2)_L$ generator. According to this condition, a $SU(2)_L$ doublet has hypercharge $Y = \pm 1/2$, a $SU(2)_L$ triplet has hypercharge $Y = 0, \pm 1$, and so on.

Note that the logic leading to the stability condition (9) made no assumption about renormalizability. Therefore, the DM candidates are absolutely stable if we strictly enforce MFV, even at the nonrenormalizable level. We also made no assumptions about the Lorentz quantum numbers of χ, so one may consider scalars, fermions, vectors, and so on. Furthermore, if additional particle content is added besides the DM candidate our stability condition holds provided we demand that χ contains the lightest mass eigenstate among the new particles.

There exist variations in the precise implementation of MFV. For example, in the case of 'linear MFV', one imagines that each Yukawa insertion comes with a small coefficient, so that only the leading terms are important, whereas in 'general MFV' all Yukawa insertions are considered to be of similar size [18]. The number of Yukawa spurions in Eq. (2) was arbitrary, so that the stability condition (9) holds for general MFV. The case of linear MFV is more restrictive and may thus lead to additional DM candidates.

Accounting for the different possible Lorentz, gauge, and flavor quantum numbers, we conclude that the MFV principle leads to a wide variety of DM candidates. MFV thus provides an attractive alternative to other well-known stabilization symmetries, such as the canonical Z_2 parity, as well as higher Abelian or non-Abelian discrete or continuous symmetries [19]. To be viable, these models must pass a number of constraints from cosmology and experiment. We now discuss these issues in an example model with a flavored DM candidate.

Flavor $SU(3)_Q$ triplet dark matter. To illustrate in more detail the general phenomenological considerations in the flavored DM scenario, here we examine in some detail the physics of a simple flavored DM candidate: a $SU(3)_Q$ triplet, SM gauge singlet scalar field. We consider an effective theory with higher dimensional operators coupling the DM to quarks, and examine the constraints coming from cosmology, direct and indirect detection probes, and cosmology. We shall see that the theory has a rich phenomenology and leads to novel phenomena at hadron colliders.

For concreteness, we consider a model with a scalar field with the following $SU(3)_c \times SU(2)_L \times U(1)_Y \times G_q$ quantum numbers:

$$S \sim (\mathbf{1}, \mathbf{1}, 0)_{\mathrm{SM}} \times (\mathbf{3}, \mathbf{1}, \mathbf{1})_{G_q}. \tag{10}$$

The Lagrangian is

$$\mathcal{L} = \partial^\mu S_i^* \partial_\mu S_i - V(S_i, H) + \mathcal{L}_{\mathrm{eff}}, \tag{11}$$

where $i = 1, 2, 3$ is a flavor index, $V(S_i, H)$ is the scalar potential involving S_i and the SM Higgs H, and $\mathcal{L}_{\mathrm{eff}}$ contains higher dimensional operators involving two S fields and two quark fields.

Spectrum. Let us first consider the scalar potential, which determines the mass spectrum of the flavored DM multiplet. The scalar S may have a bare mass term as well as contribution from electroweak symmetry breaking due to a Higgs portal coupling [16]. The relevant terms in the potential are

$$\begin{aligned} V \supset\, & m_S^2 S_i^* (a\, \mathbf{1}_{ij} + b\, (Y_u Y_u^\dagger)_{ij} + \ldots) S_j \\ & + 2\lambda\, S_i^* (a' \mathbf{1}_{ij} + b' (Y_u Y_u^\dagger)_{ij} + \ldots) S_j\, H^\dagger H, \end{aligned} \tag{12}$$

where a, b, a', b' are dimensionless MFV parameters. The ellipsis indicate further MFV spurion insertions involving Y_d which generally lead to mass fine-splittings; for simplicity we neglect these terms. The Higgs field obtains a vacuum expectation value $\langle H \rangle = v/\sqrt{2}$, breaking the electroweak gauge symmetry and giving a contribution to the S mass-squared matrix:

$$\mathcal{L} \supset -S_i^* \left[m_A^2 \mathbf{1}_{ij} + m_B^2 (Y_u Y_u^\dagger)_{ij} \right] S_j, \tag{13}$$

where we have defined

$$\begin{aligned} m_A^2 &= m_S^2 a + \lambda v^2 a', \\ m_B^2 &= m_S^2 b + \lambda v^2 b'. \end{aligned} \tag{14}$$

Without loss of generality, we freeze the Yukawa spurions to the background values $Y_d = \lambda_d$, $Y_u = V^\dagger \lambda_u$, where $(\lambda_u)_{ij} = y_{u,i} \delta_{ij}$, $(\lambda_d)_{ij} = y_{d,i} \delta_{ij}$ are diagonal in the physical Yukawa couplings $y_{u,i}$ and $y_{d,i}$; quark mass eigenstates are then obtained via $u_L \to V^\dagger u_L$. The mass matrix in Eq. (13) is diagonalized by $S \to V^\dagger S$:

$$\mathcal{L} \to -S_i^* \left[m_A^2 \mathbf{1}_{ij} + m_B^2 (\lambda_u^2)_{ij} \right] S_j. \tag{15}$$

Up to fine splittings induced by up and charm Yukawa couplings, we find $m_1^2 \simeq m_2^2 \simeq m_A^2$ and $m_3^2 \simeq m_A^2 + m_B^2 y_t^2$. There are two possibilities for the spectrum: 1) a "normal" spectrum with two lighter, nearly degenerate states $S_{1,2}$ and one heavier state S_3, or 2) an "inverted" spectrum with two heavier states $S_{1,2}$ and one lighter state S_3.

Of phenomenological importance are the trilinear couplings of scalars to the Higgs h [16], which in the mass eigenbasis read:

$$\begin{aligned} \mathcal{L} &\supset -2\lambda v h S_i^* (a' \mathbf{1}_{ij} + b' (\lambda_u^2)_{ij}) S_j \\ &\equiv -2\widetilde{\lambda}_i v h S_i^* S_i, \end{aligned} \tag{16}$$

where we have defined $\widetilde{\lambda}_i \equiv \lambda(a' + b' y_{u,i}^2)$ These couplings provide annihilation channels into SM particles, and are furthermore constrained by direct and indirect DM searches. For light DM particles in the $10 - 100$ GeV range, the recent XENON100 null results [20] imply a limit of $\widetilde{\lambda}_i \lesssim 0.015\ (m_i/10\ \mathrm{GeV})(m_h/120\,\mathrm{GeV})^2$ for the DM particle. In addition, the Higgs portal interactions above lead to nonstandard decay channels of the SM Higgs, $h \to S_i^* S_i$, provided that the scalars S_i are light enough. The partial width for $h \to S_i^* S_i$ is given by

$$\Gamma_{h \to S_i^* S_i} = \frac{\widetilde{\lambda}_i^2 v^2}{4\pi m_h} \left(1 - \frac{4 m_i^2}{m_h^2} \right)^{1/2}. \tag{17}$$

For values of $\widetilde{\lambda}_i$ on the order of the b-quark Yukawa coupling y_b, these decay modes become competitive with the SM $h \to b\bar{b}$ mode for a light Higgs.

Effective theory. By construction, the scalars do not couple to quarks at the renormalizable level since such operators lead to the decay of the would-be DM candidate. However, couplings between pairs of DM multiplets and quarks may occur in an effective theory through higher dimensional operators. At the dimension six level, we can write the following effective operators coupling two scalar multiplets to a quark and anti-quark:

$$\mathcal{O}^1_{ijk\ell} = (\bar{Q}_i \gamma^\mu Q_j)(S_k^* \overleftrightarrow{\partial_\mu} S_\ell), \tag{18}$$

$$\mathcal{O}^2_{ijk\ell} = (\bar{u}_{Ri} \gamma^\mu u_{Rj})(S_k^* \overleftrightarrow{\partial_\mu} S_\ell), \tag{19}$$

$$\mathcal{O}^3_{ijk\ell} = (\bar{d}_{Ri} \gamma^\mu d_{Rj})(S_k^* \overleftrightarrow{\partial_\mu} S_\ell), \tag{20}$$

$$\mathcal{O}^4_{ijk\ell} = (\bar{Q}_i u_{Rj})(S_k^* S_\ell) H^\dagger + \text{h.c.}, \tag{21}$$

$$\mathcal{O}^5_{ijk\ell} = (\bar{Q}_i d_{Rj})(S_k^* S_\ell) H + \text{h.c.}, \tag{22}$$

where i, j, k, l are flavor indices. From these operators we can construct the effective Lagrangian:

$$\mathcal{L}_{eff} = \frac{1}{\Lambda^2} \sum_{I=1}^{5} c^I_{ijk\ell} \mathcal{O}^I_{ijk\ell}. \tag{23}$$

The coefficients $c^I_{ijk\ell}$ in general contain all possible flavor contractions consistent with MFV. For example, for the operator $\mathcal{O}^1$ we have

$$\begin{aligned} c^1_{ijk\ell} &= c^1_1 \mathbf{1}_{ij} \mathbf{1}_{k\ell} + c^1_2 \mathbf{1}_{i\ell} \mathbf{1}_{kj} + c^1_3 (Y_u Y_u^\dagger)_{ij} \mathbf{1}_{k\ell} \\ &+ c^1_4 \mathbf{1}_{ij} (Y_u Y_u^\dagger)_{k\ell} + c^1_5 (Y_u Y_u^\dagger)_{i\ell} \mathbf{1}_{kj} \\ &+ c^{1*}_5 \mathbf{1}_{i\ell} (Y_u Y_u^\dagger)_{kj} + \ldots, \end{aligned} \tag{24}$$

where c^1_i are Wilson coefficients for a given flavor contraction, and where we display the leading terms in the MFV expansion up to one insertion of $Y_u Y_u^\dagger$.

Such higher-dimension operators can have a significant impact on the cosmology by providing the DM with efficient annihilation channels to SM quarks, as long as the effective scale $\Lambda/\sqrt{c^I_i}$ suppressing the operators is not too much larger than the weak scale. The same operators can potentially be probed by direct and indirect detection experiments, colliders, and precision flavor studies.

Indeed, certain operators and flavor structures are more constrained than others. To illustrate this fact, consider for concreteness the "inverted" spectrum from Eq. (15) in which the DM particle is S_3. For the operator $\mathcal{O}^1$, the flavor structures associated with the coefficients c^1_1 and c^1_4 in Eq. (24) lead to the vector coupling of the DM particle S_3 to valence quarks in the nucleon, and are therefore strongly constrained by direct detection experiments. On the other hand, the flavor structures associated with c^1_2, c^1_3, and c^1_5 imply that the vector couplings of the DM state S_3 are dominantly to the third generation quarks, while the couplings to valence quarks either vanish or are Yukawa and/or CKM suppressed. Therefore, these coefficients are far less constrained by direct DM searches.

Continuing on with this example, there is the possibility of new FCNCs induced by the operator $\mathcal{O}^1$ once the DM multiplet is integrated out. Whereas the coefficients c^1_1 and c^1_4 in Eq. (24) lead to strictly flavor-diagonal couplings in the quarks, the flavor structures associated with c^1_2, c^1_3, and c^1_5 indeed do induce FCNCs in the down-quark sector due to the presence of CKM mixing.

A number of other phenomenological considerations must be taken into account, each of which potentially constrain the Wilson coeffcients c^I_{ijkl} for the operators in Eqs. (19-22). Depending on the UV completion some of these coefficients may be sizable while others may be suppressed or vanish. From the effective theory perspective, one can explore the consequences of each operator on the cosmology and phenomenology in a model-independent way. Understanding which operators are constrained and which are allowed may then point towards the underlying dynamics at scales greater than Λ. A comprehensive analysis of the effective operators (18-22) and their various flavor structures will be left for future work. Instead, we shall specialize to one particular operator and examine in detail the associated cosmology and phenomenological constraints and prospects.

For the remainder of the paper let us consider operator $\mathcal{O}^5$ in Eq. (22) with the following flavor structure:

$$\mathcal{L}_{\text{eff}} = \frac{c}{\Lambda^2} [\bar{Q}_i S_i][S_j^* (Y_d)_{jk} d_{Rk}] H + \text{h.c}, \tag{25}$$

where the flavor indices are contracted within the brackets, while the Lorentz, color, and $SU(2)_L$, indices are contracted outside. The coefficient c is taken here to be a phase, so that $|c| = 1$. As we will demonstrate, this operator allows for a viable cosmology while being consistent with a variety of constraints.

After electroweak symmetry breaking and diagonalization of the quark and scalar masses, we are left with the following effective Lagrangian:

$$\mathcal{L}_{\text{eff}} \to \frac{c}{\Lambda^2} \frac{v}{\sqrt{2}} [\bar{d}_{Li} V^\dagger_{ij} S_j][S_k^* (V \lambda_d)_{k\ell} d_{R\ell}] + \text{h.c.}\,. \tag{26}$$

Focusing on the "inverted" spectrum below, $m_3 < m_1 \simeq m_2$, S_3 is the stable DM candidate, while S_1 and S_2 are unstable.

Relic abundance. First we consider the thermal relic abundance of S_3. If the DM mass m_3 is larger than the bottom quark mass m_b, the dominant annihilation mode is $S_3 S_3^* \to \bar{b} b$, coming from the following term contained in Eq. (26):

$$\mathcal{L}_{\text{eff}} \supset \frac{c}{\Lambda^2} m_b |V_{tb}|^2 S_3^* S_3 \bar{b}_L b_R + \text{h.c.}\,, \tag{27}$$

where $|V_{tb}| \simeq 1$ is the top-bottom CKM matrix element. The above term in the Lagrangian leads to the following thermally averaged annihilation cross section:

$$\begin{aligned} \langle \sigma v \rangle_{33 \to \bar{b} b} &= \frac{3}{4 \pi \Lambda^4} m_b^2 |V_{tb}|^4 \left(1 - \frac{m_b^2}{m_3^2}\right)^{1/2} \\ &\times \left\{ [\text{Re}(c)]^2 \left(1 - \frac{m_b^2}{m_3^2}\right) + [\text{Im}(c)]^2 \right\}. \end{aligned} \tag{28}$$

In the limit $m_3 \gg m_b$, the annihilation cross section approximately scales as

$$\langle \sigma v \rangle_{33 \to \bar{b}b} \simeq 1\,\mathrm{pb} \left(\frac{200\,\mathrm{GeV}}{\Lambda} \right)^4 , \tag{29}$$

suggesting that an effective scale of $\Lambda \approx 200$ GeV is required in order to match with the nominal 3σ DM abundance $\Omega^{3\sigma}_{DM} h^2 = 0.1109 \pm 0.017$ as inferred from the seven year data sample of the WMAP satellite experiment [21]. This can also be seen in Fig. 1 where the gray shaded band indicates the region in which $S_3 + S_3^*$ attain the right relic abundance. Below the threshold of $b\bar{b}$ annihilation, $m_3 \lesssim 4.2\,\mathrm{GeV}$, the most efficient channel is $S_3^* S_3 \to s\bar{b}, b\bar{s}$ with $\langle \sigma v \rangle \propto |V_{ts}|^2 |V_{tb}|^2 m_b m_s / \Lambda^4$.

CMB constraints. Residual annihilation of S_3-pairs has its strongest impact at cosmic redshifts $z \sim 1000$ at the time of recombination of electrons with hydrogen (and helium.) The electromagnetic cascades formed in the annihilation delay recombination thus affecting the temperature and polarization anisotropies of the cosmic microwave background radiation (CMB) [22, 23]. Since the rate of injected energy scales as $\langle \sigma v \rangle n_{DM}^2 m_{DM} \propto \langle \sigma v \rangle / m_{DM}$ where n_{DM} is the DM number density, the CMB limit on the annihilation cross section can conveniently, and in a rather model-independent fashion, be expressed in the form

$$(1 - f_\nu) \langle \sigma v \rangle / m_3 < r. \tag{30}$$

Here, f_ν is the fraction of energy carried away by neutrinos and we employ $r = 0.191\,\mathrm{GeV/cm^3 s^{-1}}$ as obtained in [23]. Using the PYTHIA [24] model for e^+e^- annihilations to hadrons we have verified that f_ν always remains below 10%; as a representative value we choose $f_\nu = 0.08$. The resulting lower limit on the effective scale Λ is shown by the dashed line in Fig. 1 and is strongest for small values of m_3, excluding the otherwise cosmologically viable region $m_3 \lesssim 5\,\mathrm{GeV}$.

Direct detection. The effective operator (26) also induces the elastic scattering of S_3 on nuclei. The following terms lead to spin-independent scattering:

$$\mathcal{L}_{\text{eff}} \supset \frac{\mathrm{Re}(c)}{\Lambda^2} \sum_{i=1}^{3} m_{d_i} |V_{3i}|^2 S_3^* S_3 \bar{d}_i d_i. \tag{31}$$

Though the couplings have the usual dependence on the quark masses, as for example expected in Higgs-mediated DM scattering on nuclei, the couplings to light quarks s and d are additionally CKM suppressed. This implies that only the scattering of S_3 on the b-quark content of the nucleons is relevant in constraining the scale Λ as well as the phase c of the higher-dimensional operator (26). The nucleon matrix element under question reads

$$f_{n,b} \equiv \langle n | m_b \bar{b} b | n \rangle = m_n \frac{2}{27} f_{TG}^{(n)} \simeq 0.04, \tag{32}$$



FIG. 1. Constraints: Shown are *lower* limits on the effective scale Λ as a function of m_3 for $\mathrm{Re}(c) = 1$. The dashed line is a constraint on the annihilation cross section from CMB temperature and polarization anisotropies. The solid (red) line is the exclusion bound from the recent XENON100 direct detection experiment with 100 live-days of exposure; the dotted (blue) line shows the sensitivity of a future ton-scale liquid xenon experiment. In the lower right shaded region, the effective field theory description breaks down. Within the gray shaded band, the relic density of S_3 matches the observed DM density, $\Omega_{S_3 + S_3^*} h^2 = \Omega_{DM} h^2$.

where we have used that the fact that the b-quark matrix element can be expressed in terms of the matrix element of the gluon scalar density [25], $f_{TG}^{(n)} = 1 - \sum_{u,d,s} f_{Tq}^{(n)} \simeq 0.57$ [26].

This leads to the effective spin-independent elastic DM-nucleon cross section,

$$\sigma_n = \frac{[\mathrm{Re}(c)]^2 |V_{tb}|^2 f_{n,b}^2 \, \mu_n^2}{4\pi m_3^2 \Lambda^4} \tag{33}$$

$$\simeq 3 \times 10^{-43}\,\mathrm{cm}^2 \, [\mathrm{Re}(c)]^2 \left(\frac{10\,\mathrm{GeV}}{m_3} \right)^2 \left(\frac{200\,\mathrm{GeV}}{\Lambda} \right)^4 .$$

The upper limit on σ_n from direct detection experiment can be translated into a lower limit on Λ. The most constraining data in this respect comes from the recent 100 live-day result of the XENON100 collaboration [20]. We obtain the constraint by using Yellin's maximum gap method [27] which accounts for the 3 observed events within the 48 kg fiducial detector volume and employ the results of [28] to account for detector resolution, efficiency, and acceptance. The resulting limit in the (m_{DM}, σ_n) plane is in very good agreement with the one of [20]. For the sake of exploring future sensitivity of direct detection experiments we also simulate a ton-scale liquid xenon detector assuming a raw exposure of 1 ton×year using the detector model for XENON10 of Ref. [28].

Choosing $\mathrm{Re}(c) = 1$, the XENON100 lower limit on Λ as a function of m_3 is shown by the solid (red) line in Fig. 1. As can be seen, mass values in the range $10\,\mathrm{GeV} \lesssim m_3 \lesssim 60\,\mathrm{GeV}$ are excluded. The dotted (blue) line illustrates that a future ton-scale direct detection experiment may well be sensitive to the entire electroweak-scale parameter region of m_3, excluding values up to $m_3 \lesssim 400\,\mathrm{GeV}$. It is important to note, however, that since the overall phase c of the higher dimensional operator is a free parameter, one can avoid completely the direct detection constraints by choosing $\mathrm{Re}(c) \ll 1$ while still obtaining the correct relic abundance. Finally, the the shaded region in the lower right corner of Fig. 1—subject to the condition $\Lambda < m_3/2\pi$—indicates the part of the parameter space that does not admit a perturbative UV-completion [29].

Flavor physics. The new scalars can lead to new FCNCs at the one loop level. Here we will focus on the contribution to K^0–$\bar{K}^0$ mixing. The relevant interactions from the operator (26) that can in principle induce K^0–$\bar{K}^0$ mixing are

$$\mathcal{L}_{\mathrm{eff}} \supset \sum_{i,j} \frac{c}{\Lambda^2}\, m_s V^\dagger_{1i} V_{j2} S^*_j S_i \bar{d}_L s_R + \mathrm{h.c.}\,. \tag{34}$$

Integrating out the scalars one arrives at the following Lagrangian:

$$\mathcal{L}_{sd} = C^{sd}_S (\bar{s}_R d_L)(\bar{s}_R d_L) + \mathrm{h.c.}\,, \tag{35}$$

where the Wilson coefficient is given by

$$C^{sd}_S \simeq \frac{(c^*)^2 m_s^2}{32\pi^2\Lambda^4} (V_{td}V^*_{ts})^2 F\left(\frac{m_3^2}{m_1^2}\right), \tag{36}$$

with $F(\hat{x}) = (\hat{x}+1)(\hat{x}-1)^{-1}\log\hat{x} - 2$. The function $F(\hat{x})$ takes $\mathcal{O}(1)$ values for typical values of the masses, and vanishes for $\hat{x} = 1$. To arrive at Eq. (36) we have made use of the unitarity of the CKM matrix and assumed a mass splitting predicted by MFV (15). With $\Lambda = \mathcal{O}(v)$ as dictated by the relic abundance calculation, we observe that the operator (26) leads to a contribution in Eq. (36) which is suppressed compared to the SM value by a factor $m_s^2/m_W^2 \approx 10^{-6}$. Therefore, we are comfortably below the experimental limits. Similarly, for $B_q - \bar{B}_q$ mixing, we find operators that are suppressed compared to the SM value by a factor $m_b^2/m_W^2 \approx 10^{-3}$. Furthermore, we find that no contribution to the inclusive decay $b \to s\gamma$ is induced.

Decays of heavy dark flavors. The heavier scalars S_1 and S_2 are unstable and decay via the higher dimensional operator (26). The leading operators which are unsuppressed in powers of light quark masses $m_{s,d}$ and off-diagonal CKM elements are

$$\begin{aligned}\mathcal{L}_{\mathrm{eff}} \supset\ & \frac{c}{\Lambda^2} m_b V_{tb} V^*_{cs} S^*_3 S_2 \bar{s}_L b_R \\ &+ \frac{c}{\Lambda^2} m_b V_{tb} V^*_{ud} S^*_3 S_1 \bar{d}_L b_R + \mathrm{h.c.}\,.\end{aligned} \tag{37}$$

These operators mediate the decays

$$S_2 \to S_3 s\bar{b}, \tag{38}$$

$$S_1 \to S_3 d\bar{b}. \tag{39}$$

In the limit $m_{1,2} \gg m_b, m_3$, the three-body partial decay width for $S_i \to S_3 q\bar{b}$ is approximately given by

$$\begin{aligned}\Gamma_{i\to 3qb} &\simeq \frac{|cV_{tb}V^*_{ii}|^2 m_b^2 m_i^3}{512\pi^3\Lambda^4} \\ &\simeq 10^{-5}\,\mathrm{MeV} \times \left(\frac{m_i}{25\,\mathrm{GeV}}\right)^3 \left(\frac{200\,\mathrm{GeV}}{\Lambda}\right)^4.\end{aligned} \tag{40}$$

The scalars therefore decay promptly.

LHC prospects. A novel prediction of our flavored DM scenario is the presence of heavy dark flavors. Given a production mechanism, such heavy flavors can lead to interesting signatures at the Large Hadron Collider (LHC). The effective operator (26) can in principle lead to pair production via the parton-level processes $q_i\bar{q}_j \to S_i S^*_j$. In practice, however, the production cross section for these processes is negligible due to the Yukawa suppressed couplings in Eq. (26) (this is also why the operator faces no bounds from monojet searches at the Tevatron [29]).

Here we instead focus on production of $S_{1,2}$ via the Higgs portal interaction in Eq. (13). If the new scalar states are lighter than $m_h/2$ and have a trilinear coupling to the Higgs in Eq. (16) larger than the bottom Yukawa interaction y_b, then the Higgs decays predominantly to intermediately heavy scalars $S_{1,2}$ with a partial width given by Eq. (17). These scalars subsequently decay via the three-body processes (38,39) to a bottom quark, a light quark (d or s), and the DM candidate S_3,

$$h \to S^*_1 S_1 \to b\bar{b}d\bar{d}S^*_3 S_3, \tag{41}$$

$$h \to S^*_2 S_2 \to b\bar{b}s\bar{s}S^*_3 S_3. \tag{42}$$

We now investigate the potential of the LHC ($\sqrt{s} = 14$ TeV) to discover the processes (41,42) for the benchmark scenario given in Table II. For this choice of parameters we find that the Higgs decays almost exclusively via (41,42). The most promising channel to disentangle the signal from the large SM backgrounds is expectedly the HZ production channel, where the Z boson subsequently decays to electrons or muons. One immediate reason for this is that the resulting hard leptons provide a triggerable signal. Furthermore, if the Z boson is boosted and the Higgs recoils against the leptonically decaying Z, the leptons yield a good handle to calibrate the jet energy and to measure the amount of $\not{E}_T$ in the event. Similar configurations have been used in previous analyses for

m_h	m_1	m_2	m_3	$\widetilde{\lambda}_1$	$\widetilde{\lambda}_2$	$\widetilde{\lambda}_3$	m_S^2	λ	a	a'	b	b'
120	25	25	10	0.15	0.15	0.01	-86^2	0.35	1.14	0.429	-1.10	-0.409

TABLE II. Benchmark spectrum and couplings for LHC study. The input parameters are m_1, m_2, m_3, $\widetilde{\lambda}_1$, $\widetilde{\lambda}_2$, $\widetilde{\lambda}_3$, m_S^2 and λ. This determines the Lagrangian parameters a, a', b, b'. All mass parameters are in GeV.

a boosted SM Higgs boson which decays into a bottom quark pair [30].

We simulate the signal using MadEvent [31] and the backgrounds using Herwig++ [32]. The NLO production cross sections for the backgrounds and the signal was calculated using MCFM [33]. To select a lepton we demand it to be central and sufficiently hard,

$$|y_l| < 2.5, \qquad p_{T,l} > 20 \text{ GeV}. \tag{43}$$

For the hardest lepton in the event we further require

$$p_{T,l_1} > 80 \text{ GeV}. \tag{44}$$

The muons and electrons must be isolated, *i.e.*, the hadronic transverse energy in a cone of $R = 0.3$ around the lepton has to be $E_{T_{\text{hadronic}}} < 0.1\ E_{T,l}$. We accept events with exactly two isolated leptons. The Z bosons are reconstructed by combining two oppositely charged isolated electrons or muons, requiring

$$m_Z - 10 \text{ GeV} < m_{ll} < m_Z + 10 \text{ GeV} \tag{45}$$

and $p_{T,ll} > 150$ GeV.

We group the remaining visible final state particles into "cells" of size $\Delta\eta \times \Delta\phi = 0.1 \times 0.1$. All particles in a cell are combined, and the total three-momentum is rescaled such that each cell has zero invariant mass. Cells with energy < 0.5 GeV are discarded, while the remaining ones are clustered into jets. We recombine the jets using Fastjet [34] with the anti-kT jet algorithm [35], $R = 0.7$, and

$$p_{T,j} > 30 \text{ GeV}. \tag{46}$$

We reject events where a jet with $p_{T,j} > 50$ GeV has a smaller angular separation to the Z boson than $\Delta R_{j,Z} < 1.5$. At least one jet must be present in the event, and the hardest jet in the event must be b-tagged. For the b-tag we assume an efficiency of 60% and a fake rate of 2%. The S_3 states escape the detector unobserved. Therefore we accept an event if

$$\not{E}_T > 50 \text{ GeV}, \tag{47}$$

and calculate $\not{E}_T$ from all visible objects with $|y| < 5.0$. The resulting $\not{E}_T$ vector must have a large azimuthal separation from the reconstructed Z boson,

$$\Delta\phi_{\,\not{E}_T,Z} > 2.0. \tag{48}$$



FIG. 2. Reconstruction of m_T according to Eq. (49) for a Higgs mass of 120 GeV. In addition to the cuts outlined in Eqs. (43)-(48) we require at least two jets to accept an event.

With these cuts we obtain $S/B \simeq 1$ and $S/\sqrt{B} \simeq 5$ after 15 fb^{-1}; see Table III for details. Therefore an excess in events with $\not{E}_T + b\bar{b} + l^+l^-$ can be observed relatively early at the LHC. However, due to the sizable fraction of missing transverse energy, $\not{E}_T$, the reconstruction of the mass of the Higgs boson is challenging. Because the bottom quark and the light quark from the three-body-decay of $S_{1,2}$ are highly collimated they will be observed as one jet. Thus, after requiring at least two jets in the event, in the rest frame of the Higgs one finds $m_{S_3^* S_3} \simeq m_{jj}$. Following [36], we use

$$m_T(S_i^* S_i) = \sqrt{(\not{E}'_T + E_{T,jj})^2 - (\mathbf{p}_{T,jj} + \not{\mathbf{p}}_T)^2}, \tag{49}$$

to reconstruct the Higgs mass as shown in Fig. 2. In Eq. (49), the transverse energies are given by $\not{E}'_T = \sqrt{\not{\mathbf{p}}_T^2 + m_{jj}^2}$ and $E_{T,jj} = \sqrt{\mathbf{p}_{T,jj}^2 + m_{jj}^2}$. m_{jj} is the invariant mass of the two hardest jets in the event which have a large angular separation to the Z boson, $R_{j,Z} > 2.0$. For a rough mass reconstruction of the Higgs boson in this channel we expect $\mathcal{O}(100)$ fb^{-1}, but other Higgs production channels could be added. If $m_{1,2} \lesssim m_3 + m_b$ the Higgs decays to jets and missing transverse energy. This signature displays strong similarities with the so-called buried Higgs models [37]. In either case, jet substructure can help to reconstruct the Higgs boson [38].

Discussion. In this paper we have demonstrated that the MFV hypothesis, motivated by the need to sup-

	FDM	$t\bar{t}$	ZZ	WZ	WW
$n_j \geq 1$, $n_l = 2$ and $p_{T,l_1} > 80$ GeV	12.7	8903.7	202.3	168.5	242.2
$\not{E}_T > 50$ GeV	7.8	5744.1	20.6	20.4	118.8
Z reconst. and $p_{T,Z} > 150$ GeV, no $\Delta R_{j_{50},Z} < 1.5$	4.3	9.9	5.8	3.8	0.7
$\Delta\phi_{\not{E}_T,Z} > 2.0$	4.2	4.6	5.2	3.3	0.03
b-tag	2.2	2.2	0.2	0.1	0.01

TABLE III. Cross sections for different steps in the analysis for the signal from the flavored dark matter model (FDM) and the dominant SM backgrounds. Systematic uncertainties in the measurement of $\not{E}_T$ can elevate Z+jets to a relevant background but their simulation is beyond the scope of this work. All cross sections are normalized to NLO QCD precision, calculated using MCFM, and are given in femtobarn. The background events were generated using Herwig++ and the signal was generated using MadEvent and showered using Herwig++.

press dangerous flavor changing processes in new physics scenarios, also provides a novel organizing principle for the physics of DM. Indeed, MFV leads to the stability of the lightest neutral particle for a large number of flavor representations. We have determined which representations have stable particles. This opens up the possibility for a large class of new flavored DM candidates.

Because of the connection with flavor physics, the phenomenology of these models is very rich. In addition to providing stability, MFV is a predictive framework for DM. The mass spectrum of the flavored DM multiplet is governed by the hierarchical nature of the quark Yukawa couplings. Furthermore, the pattern of couplings between the DM and the SM is prescribed by the flavor symmetries of G_q and CKM breaking. Our example study of the effective theory of a flavor $SU(3)_Q$ triplet, gauge singlet scalar DM illustrates many of the general aspects of the cosmology and phenomenology, and leads to a viable DM scenario with testable predictions at future direct detection experiments and present day colliders. In particular, the signatures of the heavy unstable dark flavors are quite novel and can be searched for at the LHC. However, it is fair to say that we have only scratched the surface of the phenomenological implications of the flavored DM scenario.

In this paper we have focused on the flavor symmetries of the quark sector, primarily because of our still incomplete knowledge about the SM neutrino sector. However, with some assumptions about how neutrinos acquire mass the MFV hypothesis can be formulated for leptons [39], and it would therefore be interesting to investigate the possibility of leptonically flavored DM. In this case, it is conceivable that heavy dark flavors may leave their imprint via decays to DM with associated leptons.

It would be interesting investigate more generally the predictions for other flavored DM multiplets, including larger flavor representations and electroweak multiplets. For the latter, typically because of the efficient annihilation to SM gauge bosons, the DM particle must be quite heavy to be in accordance with the observed cosmic abundance [17]. Multiple flavors of stable electroweak DM may instead be comparatively light since each flavor then only constitutes a portion of the DM density.

More comprehensive studies of flavored DM effective theories is certainly warranted. Much attention as of late has been given to effective field theories of DM, notably as a means to investigate the potential of hadron colliders to probe DM-quark interactions in a manner complementary to direct detection experiments [29]. In these studies, the couplings of DM to quarks were taken to be flavor universal for vector couplings and proportional to mass for scalar couplings to avoid FCNC constraints. Of course, a less restrictive possibility is to abide by the MFV principle, which allows for more general couplings between DM and quarks, including flavor violating couplings governed by the CKM elements.

Ultimately, since flavored multiplets cannot couple to quarks at the renormalizable level (else they would decay), it is interesting to construct UV completions to such effective theories of flavored DM. One may postulate additional particles that connect the DM multiplet to SM quarks, which themselves must also obey MFV. Indeed it is certainly interesting to speculate about a new sector of MFV matter, given that MFV has proven useful in addressing recent Tevatron anomalies, such as the like-sign dimuon asymmetry in B-decays [40, 41] and the top quark forward-backward asymmetry [42, 43]. Perhaps, the DM particle is contained in such an MFV sector.

The evidence for DM is one of many reasons to expect new physics, and yet the SM remains impeccable when confronted with experiment. This motivates new symmetry principles designed to forbid dangerous interactions of new particles, which may also have the dual purpose of stabilizing DM. The canonical example of such a symmetry is a discrete Z_2 parity. In this paper, we have shown that flavor symmetries together with the MFV principle provide another compelling possibility, thus pointing the way towards new avenues for physics beyond the SM and DM.

Acknowledgements

We thank Uli Nierste, Nilendra Deshpande, and Maxim Pospelov for helpful discussions. BB is grateful to Jure Zupan for stimulating discussions about his related work [44]. Research at the Perimeter Institute is supported in part by the Government of Canada through NSERC and by the Province of Ontario through MEDT. MS was supported by US Department of Energy under contract number DE-FG02-96ER40969. MS thanks the Perimeter Institute for hospitality.

---

[*] bbatell@perimeterinstitute.ca
[†] jpradler@perimeterinstitute.ca
[‡] mspannow@uoregon.edu

[1] *see* e.g. G. Jungman, M. Kamionkowski and K. Griest, Phys. Rept. **267**, 195 (1996) [arXiv:hep-ph/9506380]; G. Bertone, D. Hooper and J. Silk, Phys. Rept. **405**, 279 (2005) [arXiv:hep-ph/0404175].

[2] R. S. Chivukula and H. Georgi, Phys. Lett. B **188**, 99 (1987); L. J. Hall and L. Randall, Phys. Rev. Lett. **65**, 2939 (1990); A. J. Buras, P. Gambino, M. Gorbahn, S. Jager and L. Silvestrini, Phys. Lett. B **500**, 161 (2001) [arXiv:hep-ph/0007085].

[3] G. D'Ambrosio, G. F. Giudice, G. Isidori and A. Strumia, Nucl. Phys. B **645**, 155 (2002) [arXiv:hep-ph/0207036].

[4] E. Nikolidakis and C. Smith, Phys. Rev. D **77**, 015021 (2008) [arXiv:0710.3129 [hep-ph]].

[5] J. March-Russell, C. McCabe, M. McCullough, JHEP **1003**, 108 (2010). [arXiv:0911.4489 [hep-ph]].

[6] J. Kile and A. Soni, arXiv:1104.5239 [hep-ph].

[7] M. Hirsch, S. Morisi, E. Peinado and J. W. F. Valle, arXiv:1007.0871 [hep-ph]; D. Meloni, S. Morisi and E. Peinado, arXiv:1011.1371 [hep-ph]; M. S. Boucenna, M. Hirsch, S. Morisi, E. Peinado, M. Taoso and J. W. F. Valle, arXiv:1101.2874 [hep-ph]; D. Meloni, S. Morisi and E. Peinado, arXiv:1104.0178 [hep-ph].

[8] N. Haba, Y. Kajiyama, S. Matsumoto, H. Okada and K. Yoshioka, Phys. Lett. B **695**, 476 (2011) [arXiv:1008.4777 [hep-ph]]; Y. Kajiyama and H. Okada, Nucl. Phys. B **848**, 303 (2011) [arXiv:1011.5753 [hep-ph]]; Y. Kajiyama, H. Okada and T. Toma, arXiv:1104.0367 [hep-ph].

[9] M. Blennow, B. Dasgupta, E. Fernandez-Martinez and N. Rius, JHEP **1103**, 014 (2011) [arXiv:1009.3159 [hep-ph]].

[10] A. V. Manohar and M. B. Wise, Phys. Rev. D **74**, 035009 (2006) [arXiv:hep-ph/0606172].

[11] Y. Grossman, Y. Nir, J. Thaler, T. Volansky and J. Zupan, Phys. Rev. D **76**, 096006 (2007) [arXiv:0706.1845 [hep-ph]].

[12] C. W. Bauer, Z. Ligeti, M. Schmaltz, J. Thaler and D. G. E. Walker, Phys. Lett. B **690**, 280 (2010) [arXiv:0909.5213 [hep-ph]].

[13] J. M. Arnold, M. Pospelov, M. Trott and M. B. Wise, JHEP **1001**, 073 (2010) [arXiv:0911.2225 [hep-ph]].

[14] E. Gross, D. Grossman, Y. Nir and O. Vitells, Phys. Rev. D **81**, 055013 (2010) [arXiv:1001.2883 [hep-ph]].

[15] J. M. Arnold, B. Fornal and M. Trott, JHEP **1008**, 059 (2010) [arXiv:1005.2185 [hep-ph]].

[16] V. Silveira and A. Zee, Phys. Lett. B **161**, 136 (1985); J. McDonald, Phys. Rev. D **50**, 3637 (1994) [arXiv:hep-ph/0702143]; C. P. Burgess, M. Pospelov and T. ter Veldhuis, Nucl. Phys. B **619**, 709 (2001) [arXiv:hep-ph/0011335].

[17] M. Cirelli, N. Fornengo, A. Strumia, Nucl. Phys. **B753**, 178-194 (2006). [hep-ph/0512090].

[18] A. L. Kagan, G. Perez, T. Volansky and J. Zupan, Phys. Rev. D **80**, 076002 (2009) [arXiv:0903.1794 [hep-ph]].

[19] K. Agashe and G. Servant, Phys. Rev. Lett. **93**, 231805 (2004) [arXiv:hep-ph/0403143]; E. Ma, Phys. Lett. B **662**, 49 (2008) [arXiv:0708.3371 [hep-ph]]; T. Hambye, JHEP **0901**, 028 (2009) [arXiv:0811.0172 [hep-ph]]; F. Chen, J. M. Cline and A. R. Frey, Phys. Rev. D **80**, 083516 (2009) [arXiv:0907.4746 [hep-ph]]; D. G. E. Walker, arXiv:0907.3146 [hep-ph]; F. D'Eramo and J. Thaler, JHEP **1006**, 109 (2010) [arXiv:1003.5912 [hep-ph]]; B. Batell, Phys. Rev. D **83**, 035006 (2011) [arXiv:1007.0045 [hep-ph]]; M. Lisanti and J. G. Wacker, arXiv:0704.2816 [hep-ph]; A. Adulpravitchai, B. Batell and J. Pradler, Phys. Lett. B **700**, 207 (2011) [arXiv:1103.3053 [hep-ph]].

[20] E. Aprile *et al.* [XENON100 Collaboration], arXiv:1104.2549 [astro-ph.CO].

[21] E. Komatsu *et al.* [WMAP Collaboration], Astrophys. J. Suppl. **192**, 18 (2011) [arXiv:1001.4538 [astro-ph.CO]].

[22] T. R. Slatyer, N. Padmanabhan, D. P. Finkbeiner, Phys. Rev. **D80**, 043526 (2009). [arXiv:0906.1197 [astro-ph.CO]].

[23] G. Hutsi, J. Chluba, A. Hektor, M. Raidal, [arXiv:1103.2766 [astro-ph.CO]].

[24] T. Sjostrand, S. Mrenna, P. Z. Skands, JHEP **0605**, 026 (2006). [hep-ph/0603175].

[25] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Phys. Lett. B **78**, 443 (1978). A. I. Vainshtein, V. I. Zakharov, M. A. Shifman, Sov. Phys. Usp. **23**, 429-449 (1980).

[26] J. R. Ellis, K. A. Olive and C. Savage, Phys. Rev. D **77** (2008) 065026 [arXiv:0801.3656 [hep-ph]].

[27] S. Yellin, Phys. Rev. **D66**, 032005 (2002). [physics/0203002].

[28] P. Sorensen, JCAP **1009**, 033 (2010). [arXiv:1007.3549 [astro-ph.IM]].

[29] M. Beltran, D. Hooper, E. W. Kolb, Z. A. C. Krusberg and T. M. P. Tait, JHEP **1009**, 037 (2010) [arXiv:1002.4137 [hep-ph]]; J. Goodman, M. Ibe, A. Rajaraman, W. Shepherd, T. M. P. Tait and H. B. P. Yu, Phys. Lett. B **695**, 185 (2011) [arXiv:1005.1286 [hep-ph]]; Y. Bai, P. J. Fox and R. Harnik, JHEP **1012**, 048 (2010) [arXiv:1005.3797 [hep-ph]]; J. Goodman, M. Ibe, A. Rajaraman, W. Shepherd, T. M. P. Tait and H. B. P. Yu, Phys. Rev. D **82**, 116010 (2010) [arXiv:1008.1783 [hep-ph]]; J. F. Fortin and T. M. P. Tait, arXiv:1103.3289 [hep-ph].

[30] J. M. Butterworth, A. R. Davison, M. Rubin, G. P. Salam, Phys. Rev. Lett. **100**, 242001 (2008). [arXiv:0802.2470 [hep-ph]]; ATL-PHYS-PUB-2009-088, ATL-COM-PHYS-2009-345; T. Plehn, G. P. Salam and M. Spannowsky, Phys. Rev. Lett. **104**, 111801 (2010) [arXiv:0910.5472 [hep-ph]]; D. E. Soper, M. Spannowsky, JHEP **1008**, 029 (2010) [arXiv:1005.0417 [hep-ph]]; J. Gallicchio, J. Huth, M. Kagan, M. D. Schwartz, K. Black, B. Tweedie, JHEP **1104**, 069 (2011) [arXiv:1010.3698 [hep-ph]]; J. -H. Kim, Phys. Rev. **D83**, 011502 (2011) [arXiv:1011.1493 [hep-ph]]; D. E. Soper, M. Spannowsky, [arXiv:1102.3480 [hep-ph]].

[31] J. Alwall *et al.*, JHEP **0709**, 028 (2007).

[32] M. Bahr *et al.*, Eur. Phys. J. C **58** (2008) 639.

[33] J. M. Campbell and R. K. Ellis, Phys. Rev. D **60**, 113006 (1999); J. M. Campbell and R. K. Ellis, arXiv:1007.3492 [hep-ph]; `http://mcfm.fnal.gov/`

[34] M. Cacciari and G. P. Salam, Phys. Lett. B **641**, 57 (2006); M. Cacciari, G. P. Salam and G. Soyez, `http://fastjet.fr`

[35] M. Cacciari, G. P. Salam, G. Soyez, JHEP **0804**, 063 (2008).

[36] D. L. Rainwater, D. Zeppenfeld, Phys. Rev. **D60**, 113004 (1999). [hep-ph/9906218]; N. Kauer, T. Plehn, D. L. Rainwater, D. Zeppenfeld, Phys. Lett. **B503**, 113-120 (2001). [hep-ph/0012351].

[37] S. Chang, P. J. Fox, N. Weiner, Phys. Rev. Lett. **98**, 111802 (2007) [hep-ph/0608310]; M. Carena, T. Han, G. -Y. Huang, C. E. M. Wagner, JHEP **0804**, 092 (2008) [arXiv:0712.2466 [hep-ph]]; S. Chang, R. Dermisek, J. F. Gunion, N. Weiner, Ann. Rev. Nucl. Part. Sci. **58**, 75-98 (2008). [arXiv:0801.4554 [hep-ph]]; M. Lisanti, J. G. Wacker, Phys. Rev. **D79**, 115006 (2009) [arXiv:0903.1377 [hep-ph]]; B. Bellazzini, C. Csaki, A. Falkowski, A. Weiler, Phys. Rev. **D80**, 075008 (2009) [arXiv:0906.3026 [hep-ph]]; A. Belyaev, J. Pivarski, A. Safonov, S. Senkin, A. Tatarinov, Phys. Rev. **D81**, 075021 (2010) [arXiv:1002.1956 [hep-ph]]; A. Falkowski, J. T. Ruderman, T. Volansky and J. Zupan, JHEP **1005**, 077 (2010) [arXiv:1002.2952 [hep-ph]].

[38] G. D. Kribs, A. Martin, T. S. Roy, M. Spannowsky, Phys. Rev. **D81**, 111501 (2010) [arXiv:0912.4731 [hep-ph]]; C. -R. Chen, M. M. Nojiri, W. Sreethawong, JHEP **1011**, 012 (2010) [arXiv:1006.1151 [hep-ph]]; A. Falkowski, D. Krohn, L. -T. Wang, J. Shelton, A. Thalapillil, [arXiv:1006.1650 [hep-ph]]; G. D. Kribs, A. Martin, T. S. Roy, M. Spannowsky, Phys. Rev. **D82**, 095012 (2010). [arXiv:1006.1656 [hep-ph]]; A. Katz, M. Son, B. Tweedie, JHEP **1103**, 011 (2011). [arXiv:1010.5253 [hep-ph]]; G. D. Kribs, A. Martin, T. S. Roy, [arXiv:1012.2866 [hep-ph]].

[39] V. Cirigliano, B. Grinstein, G. Isidori and M. B. Wise, Nucl. Phys. B **728**, 121 (2005) [arXiv:hep-ph/0507001].

[40] V. M. Abazov *et al.* [D0 Collaboration], Phys. Rev. D **82**, 032001 (2010) [arXiv:1005.2757 [hep-ex]].

[41] A. J. Buras, M. V. Carlucci, S. Gori and G. Isidori, JHEP **1010**, 009 (2010) [arXiv:1005.5310 [hep-ph]]; Z. Ligeti, M. Papucci, G. Perez and J. Zupan, Phys. Rev. Lett. **105**, 131601 (2010) [arXiv:1006.0432 [hep-ph]]; B. Batell and M. Pospelov, Phys. Rev. D **82**, 054033 (2010) [arXiv:1006.2127 [hep-ph]]; A. Lenz *et al.*, Phys. Rev. D **83**, 036004 (2011) [arXiv:1008.1593 [hep-ph]]; M. Trott and M. B. Wise, JHEP **1011**, 157 (2010) [arXiv:1009.2813 [hep-ph]];

[42] V. M. Abazov *et al.* [D0 Collaboration], Phys. Rev. Lett. **100**, 142002 (2008) [arXiv:0712.0851 [hep-ex]]; T. Aaltonen *et al.* [CDF Collaboration], Phys. Rev. Lett. **101**, 202001 (2008) [arXiv:0806.2472 [hep-ex]]; T. Aaltonen *et al.* [CDF Collaboration], arXiv:1101.0034 [hep-ex].

[43] B. Grinstein, A. L. Kagan, M. Trott and J. Zupan, arXiv:1102.3374 [hep-ph]; Z. Ligeti, M. Schmaltz and G. M. Tavares, arXiv:1103.2757 [hep-ph].

[44] J. Zupan, talk at *Non-Thermal Histories of the Universe Workshop*, October 18-21, 2010, Ann Arbor, MI, USA, `http://www.umich.edu/~mctp/SciPrgPgs/events/2011/NCHU/talks/Zupan.pdf`